Last Updated: December 4, 2017 (v1.2)

Health Quality Ontario

Let's make our health system healthier

Hospital Performance Series

Pre-operative testing before low-risk surgeries

Technical Appendix



Table of Contents

Purpose of the Technical Appendix	1
Indicator Technical Specifications	1
Table 1. Percentage of endoscopy procedures with pre-operative testing	1
Table 2. Percentage of ophthalmologic surgeries with pre-operative testing	3
Table 3. Percentage of other low-risk surgeries with pre-operative testing	5
Important Data Interpretation Notes	9
Contact Information	9
References	10

Purpose of the Technical Appendix

The purpose of this Technical Appendix is to accompany Health Quality Ontario's *Hospital Performance Series* Report. It provides users with the methodological details for the pre-operative testing before low-risk surgery indicators so they can better understand the results of these measures and use the results for quality improvement activities. Below is a table of updates to the Technical Appendix:

Version	Date	Change
V 1.1	September 14, 2016	Original Technical Appendix
V 1.2	December 4, 2017	 Addition of CCI code 1.UC.58, a new code in version 2015, in 'Other low-risk surgeries' indicator Addition of CCI code 1.VK.80.GZ, a new code in version 2015, in 'Other low-risk surgeries' indicator Updated CCI code 1.OD.57 to 1.OD.57.DA in the 'Other low-risk surgeries' indicator to capture only Laparoscopic Cholecystectomy

Indicator Technical Specifications

Table 1. Percentage of endoscopy procedures with pre-operative testing

GENERAL DESCRIPTION	Indicator description	The percentage of endoscopy procedures with pre-operative • Electrocardiography (ECG)
		Chest radiography (X-ray)
	Relevance	Assess utilization of testing before low-risk procedures
	Importance	A study using Ontario data from recent years has found frequent use of pre-operative electrocardiography (ECG) and chest radiography (X-ray) among low-risk surgeries (31.0% and 10.8%, respectively). Significant variation across hospitals exists as well. (1) However, there is no evidence that conducting routine ECG or chest X-ray testing in asymptomatic patients undergoing elective low-risk surgeries improves outcomes. In fact, routine testing may lead to further unnecessary downstream testing, cancellation of surgery, and increases in patient anxiety and cost.(2-4)
	Dimension of Quality	Effectiveness
	Туре	Process
	Calculation	Denominator:
NO		Description:
ORMATIC		Number of records with eligible endoscopy procedures between April 1 and March 31 of the fiscal year.
볼		Inclusions:
JRCE		Outpatient day surgery or acute in-patient settings
DEFINTION & SOURCE INFORMATION		Ontario patients
		Aged 18 and older
		Elective admission
		 Endoscopy procedure is identified by one of the following CCI intervention codes in the first intervention code field:
		 Esophagus/Stomach

		2.NA.70.BA, 2.NA.71.BA, 2.NA.71.BP, 2.NA.71.BR, 2.NF.70.BA, 2.NF.71.BA, 2.NF.71.BP, 2.NF.71.BR • Large Bowel 2.NM.70.BA, 2.NM.71.BA, 2.NM.71.BR • The date of the intervention is on the date of admission • All procedures for patients who underwent more than one eligible procedure during the reporting period are included. Exclusions: • Records with missing institution number • Hospitals that have less than 50 eligible low-risk surgeries in the six-year reporting period, i.e. from April 1st 2011 to March 31st 2017. Numerator: Description: Number of records within the denominator with a pre-operative ECG or chest X-ray test within 60
		days prior to the index procedure.
		Inclusions:
		 The test has occurred within 60 days prior to the intervention date of the endoscopy procedure
		Tests are identified based on the following professional fee codes
		o ECG:
		Fee code: G313
		o Chest X-ray:
		Fee suffix= C
		AND
		Fee codes : X090, X091 or X092
		Method:
		Numerator/Denominator * 100
	Data sources	Canadian Institute for Health Information (CIHI) Discharge Abstract Database (DAD) for acute inpatient surgery data;
		CIHI National Ambulatory Care Reporting System (NACRS) for outpatient day surgery data;
		The Ontario Health Insurance Plan (OHIP) Claims History Database (i.e. OHIP physician service claims) for pre-operative test data; and
		The Registered Persons Database (RPDB) for patient's age calculation
	Risk adjustment, age/sex standardization	Available as crude rates
GEOGRAPHY & TIMING	Available Data Periods	Fiscal year data from 2011/12 to 2016/17
GEOGRAF & TIMING	Reporting Level	Province, LHIN, hospital, and hospital corporation for multi-site hospitals

ADDITIONAL INFORMATION	Limitations	The currently available data from administrative databases do not have the information to determine the appropriateness and the reason of the pre-operative test. Some tests might be valuable, but these cannot be differentiated from unnecessary ones. However, patients who have undergone these elective low-risk surgery generally have a low number of comorbidities. (1) With the selection of the low-risk procedures and the low-risk patient group, it is unlikely that the majority of tests were ordered to evaluate new clinical symptoms or abnormal physical findings. All tests conducted within 60 days before the index low-risk surgery are included in the
		analysis. It is possible that some tests were ordered for indications other than pre-operative testing. However, this period is generally accepted by hospitals for preoperative evaluation and has been used in previous studies. (1, 5, 6)
		No validated comprehensive "low-risk" surgical procedure list exists. However, the selection of low-risk surgery in the analysis is in line with the broad definition of "low-risk procedures" outlined in existing research (1) and guidelines on perioperative cardiac evaluation .(5,7) The majority of the procedures are minimally invasive and are performed in outpatient settings.
	Comments	N/A
	Alignment	Choosing Wisely Canada recommendations, which suggest routine pre-operative testing should be avoided for asymptomatic patients undergoing low-risk surgery. (8–10)

Table 2. Percentage of ophthalmologic surgeries with pre-operative testing

Z.	Indicator description	The percentage of ophthalmologic surgeries with pre-operative
		Electrocardiography (ECG)
		Chest radiography (X-ray)
) TIC	Relevance	Assess utilization of testing before low-risk procedures
GENERAL DESCRIPTION	Importance	A study using Ontario data from recent years has found frequent use of pre-operative electrocardiography (ECG) and chest radiography (X-ray) among low-risk surgeries (31.0% and 10.8%, respectively). Significant variation across hospitals exists as well. (1) However, there is no evidence that conducting routine ECG or chest X-ray testing in asymptomatic patients undergoing elective low-risk surgeries improves outcomes. In fact, routine testing may lead to further unnecessary downstream testing, cancellation of surgery, and increases in patient anxiety and cost.(2-4)
	Dimension of Quality	Effectiveness
	Туре	Process
	Calculation	Denominator:
		Description:
MATION		Number of records with eligible ophthalmologic surgery between April 1 and March 31 of the fiscal year.
FOR		Inclusions:
		Outpatient day surgery or acute in-patient settings
URC		Ontario patients
S		Aged 18 and older
DEFINTION & SOURCE INFORMATION		Elective admission
		 Ophthalmologic surgery is identified as one of the following CCI intervention codes in the first intervention code field:
		1.CC, 1.CD, 1.CE, 1.CF, 1.CG, 1.CH, 1.CJ, 1.CL, 1.CM, 1.CN, 1.CP, 1.CQ, 1.CR, 1.CS, 1.CT, 1.CU, 1.CV, 1.CX, 1.CZ
		The date of the intervention is on the date of admission

	1	
		 All procedures for patients who underwent more than one eligible procedure during the reporting period are included.
		Exclusions:
		Records with missing institution number
		 Hospitals that have less than 50 eligible low-risk surgeries in the six-year reporting period, i.e. from April 1st 2011 to March 31st 2017.
		Numerator: Description:
		Number of records within the denominator with a pre-operative ECG or chest X-ray test within 60 days prior to the index procedure.
		Inclusions:
		 The test has occurred within 60 days prior to the intervention date of the endoscopy procedure
		 Tests are identified based on the following professional fee codes ECG:
		Fee code: G313
		o Chest X-ray:
		Fee suffix= C
		AND
		Fee codes: X090, X091 or X092
		Method:
		Numerator/Denominator * 100
	Data sources	Canadian Institute for Health Information (CIHI) Discharge Abstract Database (DAD) for acute inpatient surgery data;
		CIHI National Ambulatory Care Reporting System (NACRS) for outpatient day surgery data; The Ontario Health Insurance Plan (OHIP) Claims History Database (i.e. OHIP physician service claims) for pre-operative test data; and
		The Registered Persons Database (RPDB) for patient's age calculation
	Risk adjustment, age/sex standardization	Available as crude rates
RAPHY ING	Available Data Periods	Fiscal year data from 2011/12 to 2016/17
GEOG & TIMI	Reporting Level	Province, LHIN, hospital, and hospital corporation for multi-site hospitals
ADDITIONAL INFORMATION & TIMING	Limitations	 The currently available data from administrative databases do not have the information to determine the appropriateness and the reason of the pre-operative test. Some tests might be valuable, but these cannot be differentiated from unnecessary ones. However, patients who have undergone these elective low-risk surgery generally have a low number of comorbidities. (1) With the selection of the low-risk procedures and the low-risk patient group, it is unlikely that the majority of tests were ordered to evaluate new clinical symptoms or abnormal physical findings. All tests conducted within 60 days before the index low-risk surgery are included in the
ADE		analysis. It is possible that some tests were ordered for indications other than pre-operative

	 testing. However, this period is generally accepted by hospitals for preoperative evaluation and has been used in previous studies. (1, 5, 6) No validated comprehensive "low-risk" surgical procedure list exists. However, the selection of low-risk surgery in the analysis is in line with the broad definition of "low-risk procedures" outlined in existing research (1) and guidelines on perioperative cardiac evaluation .(5,7) The majority of the procedures are minimally invasive and are performed in outpatient settings.
Comments	N/A
Alignment	Choosing Wisely Canada recommendations, which suggest routine pre-operative testing should be avoided for asymptomatic patients undergoing low-risk surgery. (8–10)

Table 3. Percentage of other low-risk surgeries with pre-operative testing

	Indicator description	The percentage of other low-risk surgeries with pre-operative
NO	maicator acsorption	Electrocardiography (ECG)
		Chest radiography (X-ray)
	D. I	
L H	Relevance	Assess utilization of testing before low-risk procedures
GENERAL DESCRIPTION	Importance	A study using Ontario data from recent years has found frequent use of pre-operative electrocardiography (ECG) and chest radiography (X-ray) among low-risk surgeries (31.0% and 10.8%, respectively). Significant variation across hospitals exists as well. (1) However, there is no evidence that conducting routine ECG or chest X-ray testing in asymptomatic patients undergoing elective low-risk surgeries improves outcomes. In fact, routine testing may lead to further unnecessary downstream testing, cancellation of surgery, and increases in patient anxiety and cost.(2-4)
	Dimension of Quality	Effectiveness
	Туре	Process
	Calculation	Denominator:
		Description:
		Number of records with eligible low-risk surgery between April 1 and March 31 of the fiscal year.
		Inclusions:
ATIC		Outpatient day surgery or acute in-patient settings
, M		Ontario patients
SH SH		Aged 18 and older
_ 		Elective admission
DEFINTION & SOURCE INFORMATION		 Other low-risk procedure is identified as one of the following CCI intervention codes in the first intervention code field:
« ∠		o Orthopedic
<u></u> 은		 Shoulder (endoscopic drainage/extraction/procurement/release)
DEFIN		1.TA.52.DA, 1.TA.58.DA, 1.TA.72.DA
		 Clavicle (endoscopic drainage/distal resection)
		1.TB.52.GB, 1.TB.52.GD, 1.TB.87.DA
		 Rotator Cuff (endoscopic extraction/destruction/release/repair)
		1.TC.57.DA, 1.TC.59.DA 1.TC.72.DA, 1.TC.80.DA, 1.TC.80.GC
		 Arm/Forearm (Nerve decompression/repair/excision)

```
1.BM.72, 1.BM.80, 1.BM.87, 1.BN.72
```

- Wrist/Hand
- 1.UB.52, 1.UB.53, 1.UB.55, 1.UB.58, 1.UB.72, 1.UB.73, 1.UB.74, 1.UB.75,
- 1.UB.80, 1.UB.87, 1.UC.53, 1.UC.55, 1.UC.58, 1.UC.72, 1.UC.73,
- 1.UC.74, 1.UC.75, 1.UC.79, 1.UC.80, 1.UC.82, 1.UC.87, 1.UC.89,
- 1.UF.55, 1.UF.73, 1.UF.74, 1.UF.80, 1.UF.87, 1.UG.52, 1.UG.53, 1.UG.55,
- 1.UG.72, 1.UG.73, 1.UG.74, 1.UG.75, 1.UG.80, 1.UG.87, 1.UJ.71,
- 1.UJ.73, 1.UJ.74, 1.UJ.75, 1.UJ.82, 1.UJ.87, 1.UJ.93, 1.UK.53, 1.UK.55,
- 1.UK.72, 1.UK.73, 1.UK.74, 1.UK.75, 1.UK.80, 1.UK.87, 1.UK.93, 1.US.58,
- 1.US.72, 1.US.80, 1.UT.53, 1.UT.55, 1.UT.72, 1.UT.80, 1.UT.84, 1.UU.53,
- 1.UU.55, 1.UU.72, 1.UU.80, 1.UU.84, 1.UV.72, 1.UV.80, 1.UY.52,
- 1.UY.55, 1.UY.56, 1.UY.57, 1.UY.59, 1.UY.72, 1.UY.80, 1.UY.87
 - Nerve
- 1.BP.72, 1.BP.80, 1.BP.87, 1.BQ.72, 1.BQ.80, 1.BQ.87
 - Hip Arthroscopy (extraction/procurement/release/partial excision)

1.VA.58.DA, 1.VA.72.DA, 1.VA.87.GB

- Knee
 - Arthroscopy (drainage/extraction/procurement/release/partial excision)
- 1.VG.52.DA, 1.VG.58.DA, 1.VG.72.DA, 1.VG.87.GB
 - Meniscus (endoscopic repair/partial or total excision)
- 1.VK.80.DA, 1.VK.80.GZ, 1.VK.87.DA, 1.VK.89.DA
 - Ligament (ACL)(endoscopic repair/partial excision)
- 1.VL.80.DA, 1.VL.80.FY, 1.VL.87.DA, 1.VL.87.GB
- Ankle/Foot Arthroscopy (extraction/procurement/release)
- 1.WA.58.DA, 1.WA.72.DA
 - Microdiscectomy
- 1.SE.87
- Urologic
 - Bladder neck suspension
 - 1.PL.74
 - Bladder neck transurethral partial excision
 - 1.PL.87
 - Bladder Drainage
 - 1.PM.52, 1.PM.54
 - Bladder
 - 1.PM.59
 - Prostate resection (TURP)
 - 1.QT.87
 - Urethra
 - 1.PQ.26, 1.PQ.35, 1.PQ.50, 1.PQ.52, 1.PQ.53, 1.PQ.54, 1.PQ.55, 1.PQ.57, 1.PQ.58, 1.PQ.59, 1.PQ.72, 1.PQ.77, 1.PQ.78, 1.PQ.80
- Gynecologic
 - Hysteroscopy (endometrial ablation)
 - 1.RM.59.BA
 - Laparoscopy (oophorectomy, cystectomy, fixation)

1.RB.52.BA, 1.RB.52.DA, 1.RB.56.DA, 1.RB.74.DA, 1.RB.87.DA, 1.RB.89.DA, 1.RD.52.BA, 1.RD.89.DA General • Hernia repair (repair muscles of chest and abdomen) 1.SY.80 Inguinal lymph nodes 1.MJ.52, 1.MJ.87, 1.MJ.89 Peripheral lymph nodes 1.MK.52, 1.MK.87, 1.MK.89 Breast (removal of device/fixation/size reduction/size increase/repair/partial or total excision) 1.YM.55, 1.YM.74, 1.YM.78, 1.YM.79, 1.YM.80, 1.YM.87, 1.YM.89 Laparoscopic Cholecystectomy 1.OD.57.DA The date of the intervention is on the date of admission All procedures for patients who underwent more than one eligible procedure during the reporting period are included. Exclusions: Records with missing institution number Hospitals that have less than 50 eligible low-risk surgeries in the six-year reporting period, i.e. from April 1st 2011 to March 31st 2017. **Numerator:** Description: Number of records within the denominator with a pre-operative ECG or chest X-ray test within 60 days prior to the index procedure. Inclusions: The test has occurred within 60 days prior to the intervention date of the endoscopy procedure Tests are identified based on the following professional fee codes o ECG: Fee code: G313 Chest X-ray: Fee suffix= C AND Fee codes: X090, X091 or X092 Method: Numerator/Denominator * 100 Data sources Canadian Institute for Health Information (CIHI) Discharge Abstract Database (DAD) for acute inpatient surgery data;

Hospital Performance Series

claims) for pre-operative test data; and

CIHI National Ambulatory Care Reporting System (NACRS) for outpatient day surgery data; The Ontario Health Insurance Plan (OHIP) Claims History Database (i.e. OHIP physician service

		The Registered Persons Database (RPDB) for patient's age calculation
	Risk adjustment, age/sex standardization	Available as crude rates
GEOGRAPHY & TIMING	Available Data Periods	Fiscal year data from 2011/12 to 2016/17
GEOG & TIM	Reporting Level	Province, LHIN, hospital, and hospital corporation for multi-site hospitals
ADDITIONAL INFORMATION GEOG & TIM	Limitations	 The currently available data from administrative databases do not have the information to determine the appropriateness and the reason of the pre-operative test. Some tests might be valuable, but these cannot be differentiated from unnecessary ones. However, patients who have undergone these elective low-risk surgery generally have a low number of comorbidities. (1) With the selection of the low-risk procedures and the low-risk patient group, it is unlikely that the majority of tests were ordered to evaluate new clinical symptoms or abnormal physical findings. All tests conducted within 60 days before the index low-risk surgery are included in the analysis. It is possible that some tests were ordered for indications other than pre-operative testing. However, this period is generally accepted by hospitals for preoperative evaluation and has been used in previous studies. (1, 5, 6) No validated comprehensive "low-risk" surgical procedure list exists. However, the selection of low-risk surgery in the analysis is in line with the broad definition of "low-risk procedures" outlined in existing research (1) and guidelines on perioperative cardiac evaluation .(5,7) The majority of the procedures are minimally invasive and are performed in outpatient settings.
	Comments	N/A
	Alignment	Choosing Wisely Canada recommendations, which suggest routine pre-operative testing should be avoided for asymptomatic patients undergoing low-risk surgery. (8–10)

Important Data Interpretation Notes

Data suppression due to privacy: to ensure privacy, when numerators and/or denominators are between 1 and 5, all values, including numerator, denominator and rate are suppressed and denoted by the symbol "†". Please note that in order to avoid back calculation, suppressed hospital site level data are not included in the hospital corporation level reporting.

No procedures during reporting period: all values, including numerator, denominator and rate are denoted by the symbol "§" if no selected low-risk surgical procedures were performed within the reporting period.

Flag for unstable rates: indicator rates are considered as unstable and are flagged with an asterisk "*" if the denominator is between 6 and 29. In this case, the data should be interpreted with caution.

Data used for the institutional variation graph/analysis: The most recent fiscal year, i.e., FY2016/17 data are used in the institutional variation bar graphs and corresponding range analysis. Please note that hospitals with suppressed data or without any selected low-risk procedures are not included. Hospitals with flagged unstable rates are included.

Data distribution analysis in the institutional variation section: In order to help hospitals better understand their performance compared to others, the following distribution data are provided in a table under the institutional variation graph: minimum, maximum, and quartiles (i.e., the 25th percentile, median, and the 75th percentile). These statistics are defined as follows:

- **Minimum and maximum**: Upon ordering a set of numbers from smallest to largest, the smallest number is the *minimum* and the largest number is the *maximum*.
- **Quartiles**: The quartiles of a ranked set of numbers are the three points that divide the data set into four equal groups, each group comprising a quarter of the data. It includes the 25th percentile, median and the 75th percentile. The 25th percentile, i.e., the first quartile, is defined as the middle number between the minimum and the median. The *median* is the second quartile, and is the middle number in the ranked set. The 75th percentile is the middle value between the median and the maximum.

Contact Information

Health Quality Ontario 130 Bloor Street West, 10th floor Toronto, ON M5S 1N5

Telephone: 416-323-6868 **Toll-free:** 1-866-623-6868

Fax: 416-323-9261

Email: HospitalReport@hqontario.ca

Website: www.hgontario.ca

References

- 1. Kirkham KR, Wijeysundera DN, Pendrith C, Ng R, Tu JV, Laupacis A, et al. Preoperative testing before low-risk surgical procedures. CMAJ. 2015; 187(11): E349–E358
- 2. Chung F, Yuan H, Yin L, Vairavanathan S, Wong DT. Elimination of preoperative testing in ambulatory surgery. *Anesth Analg.* 2009 Feb 1;108(2):467-475.
- 3. Fritsch G, Flamm M, Hepner DL, Panisch S, Seer J, Soennichsen A. Abnormal pre-operative tests, pathologic findings of medical history, and their predictive value for perioperative complications. Acta Anaesthesiol Scand. 2012 Mar 1;56(3):339-350.
- 4. Institute of Health Economics. Routine preoperative tests are they necessary? [Internet]. Edmonton (AB): Institute of Health Economics; 2007 [cited 2016 June21]. Available from: http://www.ihe.ca/download/routine_preoperative_tests_are_they_necessary.pdf.
- 5. Fleisher LA, Beckman JA, Brown KA, et al. ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery: a report of the American College of Cardiology/ American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery) developed in collaboration with the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, and Society for Vascular Surgery [published erratum in *J Am Coll Cardiol* 2008;52:793-4]. *J Am Coll Cardiol* 2007; 50:e159-241.
- 6. Bugar JM, Ghali WA, Lemaire JB, Quan H. Canadian Perioperative Research Network. Utilization of a preoperative assessment clinic in a tertiary care centre. Clin Invest Med 2002; 25: 11-18.
- Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B. et al. 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. 2014; 64(22): e77-e137.
- 8. Canadian Anesthesiologists' Society. Anesthesiology: Five things physicians and patients should question [Internet]. Toronto (ON): Choosing Wisely Canada; 2015 [cited 2016 June 23]. Available from: http://www.choosingwiselycanada.org/recommendations/anesthesiology/.
- 9. Canadian Association of General Surgeons. Six things physicians and patients should question [Internet]. Toronto (ON): Choosing Wisely Canada; 2014 [cited 2016 June 23]. Available from: www.choosingwiselycanada.org/recommendations/general-surgery/.
- 10. Canadian Society of Internal Medicine. Five things physicians and patients should question [Internet]. Toronto (ON): Choosing Wisely Canada; 2014 [cited 2016 June 23]. Available from: www.choosingwiselycanada.org/recommendations /internal-medicine/